

Pumpkins: (watermelon, cantaloupe, pumpkin, ...)



0.3 – 0.4 l/t seed

Seed treatment at least 2 hours before sowing. Consumption of working solution 10-20 L/t.

15 - 30 ml/ 10 - 12 l of water

Immersion of seeds before sowing for 6-12 hours.

0.5–0.6 l/ha foliar

Spraying during the growing season 2-3 times with an interval of 10-14 days between treatments.

Main phenophases: 2-3 true leaves, budding - the beginning of flowering and the period of fruit formation (fruit growth).

2 – 2.5 l/ 1000 l of water

Watering the roots 2-4 times during the growing season with an interval of 10-14 days

Consumption of the working solution according to the irrigation rate.

The main phenophases in the open field: 2-3 true leaves, budding - the beginning of flowering and peri- from fruit formation (fruit growth).

In protected soil: 1. watering when planting seedlings / 2. watering 14 days after planting, before flowering / 3. watering 14 days after the previous one from the beginning of fruit planting / 4. 10-14 days after the previous one.

When growing plants in protected soil, for example on melons, the application of Relikt R accelerates the growth of plants and fruits. Increased nutrition can lead to cracking. When the first signs appear, it is necessary to change the rate of irrigation, reducing the amount of water by 10-20% of the usual. In this case, it should be carefully monitored so that the leaves of the culture do not start to descend too much.